

PROCARE[®]

LANDSCAPE & TURF PRODUCTS

Insecticide & Termiticide

CONCENTRATE

10% PERMETHRIN

- Kills over 100 named insect pests
- Kills troublesome outdoor insects which destroy lawns, roses, flowers, trees and shrubs, vegetables, fruits, and nuts
- Provides contact and residual control for up to 4 weeks

NET 1.0 GAL (3.79 L)

ACTIVE INGREDIENT: By Wt.
Permethrin 10.0%
INERT INGREDIENTS: 90.0%
TOTAL 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Read Entire Label and Pamphlet. Use Strictly in
Accordance with Precautionary Statements and Directions
and with Applicable State and Federal Regulations.

5.5"

5.5"

BOUND BOOKLET

For visual reference only

CONSTRUCTION



(Underneath Booklet)

.562"

4.375"

.562"

5.25"

5.5"

5.5"

LANDSCAPE & TURF PRODUCTS

10% PERMETHRIN

- Protects your lawns, trees & shrubs, roses & flowers, fruits & nuts, and vegetables from: alfalfa loopers, ants, aphids, armyworms, bagworms, beetles, borers, brown dog ticks, budworms, cabbageworms, chinch bugs, cicadas, citrus black flies, cluster flies, codling moths, corn earworms, crickets, cutworms, earwigs, elm spanworms, fall cankerworms, fire ants, fleas, green fruitworms, gnats, grasshoppers, houseflies, inchworms, Japanese beetles, leafhoppers, leafminers, leafrollers, lone star ticks, lygus bugs, mealybugs, millipedes, mole crickets, mosquitoes, moths, navel orangeworms, scales, sod webworms, sowbugs (pillbugs), spider mites, spring cankerworms, tent caterpillars, thrips, tobacco budworms, weevils, whiteflies and many others.
 - For use around foundations, outside buildings, porches, wood piles, wooden fences, decks, and patios
 - Subterranean termite control
 - Provides contact and residual control for up to 4 weeks
- 

Manufactured by:
GRO TEC, INC.
P.O. BOX 290
MADISON, GA 30650

EPA Registration No. 53883-69-59144
EPA Establishment No. 59144-GA-1



BOOKLET OUTSIDE

BACK COVER

(Last Page)

4.25"

FRONT COVER

(Page 1)

4.50"

PRECAUTIONARY STATEMENTS (continued)

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Keep Out of Reach of Children

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms including fish and invertebrates. Do not apply directly to water. Avoid contamination of ornamental fish ponds. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not use in undiluted form.

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS.

WARRANTY STATEMENT

Gro Tec, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Gro Tec, Inc. To the extent consistent with applicable law, Gro Tec, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except, as expressly provided herein, Gro Tec, Inc. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, the exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Gro Tec, Inc. election, the replacement of this product.

NOTES

PROCARE®

LANDSCAPE & TURF PRODUCTS

Insecticide & Termiticide

10% PERMETHRIN

- Protects your lawns, trees & shrubs, roses & flowers, fruits & nuts, and vegetables from: alfalfa loopers, ants, aphids, armyworms, bagworms, beetles, borers, brown dog ticks, budworms, cabbageworms, chinch bugs, cicadas, citrus black flies, cluster flies, codling moths, corn earworms, crickets, cutworms, earwigs, elm spanworms, fall cankerworms, fire ants, fleas, green fruitworms, gnats, grasshoppers, houseflies, inchworms, Japanese beetles, leafhoppers, leafminers, leafrollers, lone star ticks, lygus bugs, mealybugs, millipedes, mole crickets, mosquitoes, moths, navel orangeworms, scales, sod webworms, sowbugs (pillbugs), spider mites, spring cankerworms, tent caterpillars, thrips, tobacco budworms, weevils, whiteflies and many others.
- For use around foundations, outside buildings, porches, wood piles, wooden fences, decks, and patios
- Subterranean termite control
- Provides contact and residual control for up to 4 weeks

Manufactured by:
GRO TEC, INC.
P.O. BOX 290
MADISON, GA 30650

EPA Registration No. 53883-69-59144
EPA Establishment No. 59144-GA-1



OPEN HERE

4.75"

4.75"

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages is forty (40).**

INSIDE TEXT
(Even Pages)

BOOKLET INSIDE

INSIDE TEXT
(Odd Pages)

4.25"

4.50"

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

GENERAL PRECAUTIONS AND USE RESTRICTIONS

- Apply this product only as specified on this label.
- Do not use spoons or measuring cups for food purposes after use with pesticides
- Do not use in food areas of restaurants or food handling establishments or other areas where food is commercially prepared or processed. Do not use in a facility that is in operation or in serving areas while food is exposed. Serving areas are defined as areas where prepared foods are served such as dining rooms, but exclude areas where food may be prepared or held. When used in the home, cover or remove all exposed food and cooking utensils and cover all food-processing surfaces or wash thoroughly after treatment and before re-use.
- Do not contaminate food or feedstuffs.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying through outdoor residential misting systems, do not enter or allow others to enter until vapors, mists and aerosols have dispersed and the area has been thoroughly ventilated.
- Except when applying through outdoor residential misting systems, do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying through outdoor residential misting systems, only protected handlers may be in the area during application.
- Except when applying through outdoor residential misting systems, do not enter or allow others to enter until vapors, mists, and aerosols have dispersed, and the area has been thoroughly ventilated.
- Do not remain in the treated area. Exit immediately and remain outside the treated area until vapors, mists and aerosols have dispersed.
- Remove or cover exposed food and drinking water before application.
- Do not apply directly to or near water, storm drains or drainage ditches. Do not apply in windy conditions. To prevent run-off, do not water the treated area(s) or apply when heavy rain is expected. Rinse application equipment only over treated area.
- Do not make applications during rain.
- Do not wet plants to point of run-off or drip. Do not use treated articles until spray has dried.
- When used in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- Remove pets and birds, and cover fish aquaria before application. Turn off aquarium filtering systems before spraying and wait approximately 15 minutes after application before turning the filter back on.
- Except for outdoor residential misting applications, all outdoor applications must be limited to spot or crack-and-crevice applications, except for the following permitted uses: 1) Treatment to soil or vegetation around structures; 2) Applications to lawns, turf, and other vegetation; Application to building foundations up to a maximum height of 3 feet. Other than applications to building foundations all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches, and structural surfaces (such as windows, doors and eaves) are limited to crack and crevice applications only.

DIRECTIONS FOR USE (continued)

THREE STEPS TO MIX AND APPLY:

SHAKE WELL BEFORE USING

1. For lawns and vegetables, measure the area to be sprayed. Calculate square feet by multiplying length by width. For all other applications, apply as a thorough cover spray.
2. Use a clean sprayer. Carefully measure and mix the amount of concentrate and water as directed.
3. Spray as directed. Thorough coverage is important. Flush sprayer with clean water after each use. Make applications when insects first appear. Repeat applications as directed.

Volume conversions: 1 fl.oz. = 6 tsp. or 2 Tbsp., 8 fl.oz. = ½ pt. or 1 cup

DIRECTIONS FOR APPLYING

HOME LAWNS

TURF VARIETY	PESTS	HOW TO USE
bentgrass, bermudagrass, bluegrass, dichondra, fescue, Irish moss, merion, St. Augustine grass	ants, armyworms, brown dog ticks, chinch bugs, crickets, cutworms, fleas, grasshoppers Japanese beetle grubs, mole crickets, sod webworms, ticks (including those that carry Lyme disease), leathoppers	Thoroughly wet down grass a few hours before applying. Home lawns should be no taller than 3 inches at time of application. Slightly more water may be used as long as the amount of product per area is as listed below. For ants, crickets and grasshoppers, use ½ fl.oz. in 3 gals. of water to cover 1,000 sq.ft. For other listed insects, use 1½ fl.oz. in 10 gal. of water to cover 1,000 sq.ft.

HOUSE PLANTS INDOORS: TO PROTECT PLANTS IN RESIDENTIAL GREENHOUSES AND FOR INDOOR LANDSCAPING:

HOUSE PLANTS	PESTS	HOW TO USE
ageratum, ardisia, aster, baby's breath, begonia, birdsnest fern, bleeding heart, carnation, chrysanthemum, coleus, crown of thorns, cockscomb, cyclamen, dracaena, dumbcane, english ivy, fuchsia, gladiolus, grape ivy, juniper, lily, marigold, orchid, palm, pansy, peperomia, petunia, philodendron, piggy-back plant, poinsettia, portulaca, pothos, prayer plant, purple passion, rabbits foot fern, rose, schefflera, snapdragon, snake plant, statice, velvet plant, verbena, zinnia	aphids, armyworms, corn earworms, exposed thrips, fall cankerworms, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scales, soldier mites, spring cankerworms, whiteflies	Mix ¼ fl.oz. in 1 gal. of water. Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Use in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. As with all plants listed, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore, a wide variety of susceptibility to injury. To ascertain possible phytotoxicity, treat a few plants as described above, wait 24 hours, and check for possible injury.
velvet plants	fungus gnats	

FIRE ANT CONTROL

INDIVIDUAL MOUND AND PLANT CONTAINER TREATMENT:

Mix ½ fl.oz. in 3 gals. of water and apply as a drench using a sprinkler can. Gently apply 1 gallon of the mixture over the surface of each mound. Thoroughly wet mound and surrounding area of 3x4 ft. (12 sq.ft.). For best results apply in cool weather 650-800F early or late in the day. Treat new mounds as they appear. Pressurized sprays may disturb ants and cause migration, which will reduce product effectiveness. Use the same method as above for the control of Fire Ants that may nest in permanently placed containerized ornamentals plants.

4.75"

4.75"

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages is forty (40).**

INSIDE TEXT (Even Pages)

4.25"

BOOKLET INSIDE

INSIDE TEXT (Odd Pages)

4.50"

DIRECTIONS FOR APPLYING (continued)

TREES & SHRUBS

TREE AND SHRUB VARIETY	PESTS	HOW TO USE
Arizona, cypress, azalea, birch, cherry, non-bearing citrus, conifers, elm, English ivy, euonymus, weeping fig, fir, honeysuckle, lilac, mock-orange, oak, palm, philodendron, pine, poinsettia, tulip poplar, rhododendron, taxus	aphids, bagworms, cicadas, exposed thrips, fall cankerworms, inchworms, leafminers, leafrollers, mealybugs, spider mites, whiteflies, gypsy moth, tent caterpillar, Japanese beetles	Use ¼ fl.oz. in 1 gal. of water. Apply as a thorough spray, wetting leaves and branches to the dripping point. Try to penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50°-75°F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.
cherry, honeysuckle	fall webworms	
pine	pine beetles, pine moths, pine needle miners, needle scales	
other conifers	budworms	
Douglas and grand firs	tussock moth	
non-bearing citrus	citrus black flies	
oak	oakworms	
elm	elm leaf beetles, elm spanworms, fall webworms	

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. Wet tents with spray on all sides. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

ROSES & FLOWERS (ORNAMENTALS)

TREE AND SHRUB VARIETY	PESTS	HOW TO USE
ageratum, aster, ardisia, azalea, baby's breath, begonia, birds-nest fern, bleeding heart, carnation, chrysanthemum, coleus, common ninebark and snowberry, crown of thorns, cockscomb, cyclamen, dracaena, dumbcane, english ivy, exacum, fuchsia, gladiolus, grape ivy, gold bells, hypoestes, ivy, juniper, lilac, marigold, mock-orange, nannysberry, orchid, palm, pansy, pea shrub, peperomia, petunia, philodendron, piggy-back plant, poinsettia, portulaca, pothos, prayer plant, purple passion, rabbits foot fern, rhododendron, rose, schefflera, snapdragon, snake plant, statice velvet plant, verbenia, zinnia	aphids, armyworms, bagworms, cabbage worms, cicadas, cornworms, cutworms, exposed thrips, fall cankerworms, fall webworms, fungus gnat, inchworms, Japanese beetles, leafminers, leafrollers, mealybugs, scale crawlers, spring cankerworms, spider mites, whiteflies, rose chafer, psyllids, scale insects	Use ¼ fl.oz. in 1 gal. of water. Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the later afternoon or evening when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Application can be made up to the day of collection.

FRUIT & NUT TREES

GENERAL DIRECTIONS: Use ½ fl.oz. in 1 gal. of water. Apply when insects appear. Wet the plants to the dripping point. Cover undersides of leaves and penetrate dense foliage to the extent possible.

TREE VARIETY	PESTS	HOW TO USE
almonds, pecans	navel orangeworms, peach twig borers	Do not make more than 2 applications during hull split and 5 applications per season. Can be applied up to 7 days prior to harvest.
apples	green fruitworms, aphids, leafrollers (oblique-banded, redbanded), plum curculio, rosy apple aphids, spotted tentiform leafminers, tarnished plant bugs, white apple leafhoppers, Japanese beetles	Do not make more than 3 applications. Do not apply after petal fall.

4.75"

DIRECTIONS FOR APPLYING (continued)

FRUIT & NUT TREES (continued)

TREE VARIETY	PESTS	HOW TO USE
peaches	borers (lesser peachtree, peach twig), green fruitworms, plum curculio, Oriental fruit moths, tarnished plant bugs	Do not apply within 7 days of harvest. Do not make more than 8 applications per season.
pears (dormant through delayed dormant)	pear psylla	Apply during the dormant through delayed dormant growth period only. Do not make more than 2 applications per season.
pears (summer)	codling moths, green fruitworms, pear psylla, aphids	Do not make more than 3 applications per season. Can be applied up to 14 days prior to harvest.

VEGETABLES

Prepare a spray solution at the dilution rates listed for each crop in the chart below. Apply the spray solution at the rate of 1 gallon to cover 440 square feet surface area.

VEGETABLES	PESTS	HOW TO USE
asparagus (pre-harvest)	asparagus beetles, cutworms	Use ¾ fl.oz. in 2 gals. of water. Apply to ensure uniform coverage. Do not apply more than 4 applications per crop. Can be applied up to 3 days prior to harvest.
asparagus (post-harvest)	larval stages of asparagus beetle; tarnished plant bugs, lygus bugs, adult Japanese beetle	Use ¼ fl.oz. in 1 gal. of water. Treat ferns or bush growth after spear harvest when insects are present.
celery - see leafy vegetables		
cole crops, broccoli, brussels sprouts, cabbage, cauliflower	beet armyworms, cabbage aphids and loopers, diamondback moths, imported cabbageworms	Use ¼ fl.oz. in 1 gal. of water. Apply every 5 days or as needed. Can be applied up to 1 day prior to harvest. Do not make more than 8 applications on cauliflower, Brussels sprouts and broccoli and 10 applications on cabbage.
corn - see sweet corn		
cucumber, melons, pumpkins, squash	flea beetle, leafhopper	Use ½ fl.oz. in 1 gal. water. Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.
cucurbits: balsam pear (bitter melon), Chinese waxgourd (Chinese preserving melon), citron melon, cucumber, gherkin, edible gourds; melons including hybrids such as cantaloupe, casaba, crenshaw, honeydew, honey balls, mango melon, muskmelon, Persian melon, pumpkin, summer & winter squash, watermelon	aphids, cabbage looper, cucumber beetle (adults), cutworms, leafhoppers, leafminers, melonworms, pickleworm, plant bugs, squash bugs.	Use ½ fl.oz. in 1 gal. water. Apply to ensure uniform coverage. Can be applied up to the day of harvest.
eggplant	Colorado potato beetles, vegetable leafminers	Use ¾ fl.oz. in 1 gal. of water. Apply to ensure uniform coverage. Do not exceed 16 applications. Can be applied up to 3 days prior to harvest.
horseradish (foliar application)	imported crucifer weevils	Use ¾ fl.oz. in 1 gal. water. Do not exceed 3 foliar applications. Can be applied up to 22 days prior to harvest.

4.75"

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages is forty (40).**

INSIDE TEXT (Even Pages)

BOOKLET INSIDE

INSIDE TEXT (Odd Pages)

4.25"

4.50"

DIRECTIONS FOR APPLYING (continued)

VEGETABLES (continued)

VEGETABLES	PESTS	HOW TO USE
head lettuce	corn earworms, beet armyworms, loopers (alfalfa, cabbage), tobacco budworms	Use ½ fl.oz. in 1 gal. water. Apply every 5-10 days as needed. Do not make more than 10 applications per season. Can be applied up to 1 day prior to harvest.
leafy vegetables: amaranth, arrugula, celery, celituce, chervil, corn salad, chrysanthemum (edible-leaved and garland), cress (garden and upland), dandelion, dock, endive, fennel, lettuce(head and leaf), okra, parsley, purslane (garden and winter), rhubarb, spinach, Swiss chard	alfalfa looper, aphids, beet armyworm, cabbage looper, corn earworm, cutworms, european corn borer, fall armyworm, green cloverworm, southern armyworm, tobacco budworm, vegetable leafminer.	Use 1/3 fl.oz. in 1 gal. water. Apply every 5 to 10 days as needed. Can be applied up to 1 day prior to harvest.
melons-see cucurbits		
potatoes	beet armyworm, Colorado potato beetle, potato leafhopper, potato tuberworm, cabbage looper, cutworms, European corn borer, potato psyllid, tarnished plant bug	Use ¾ fl.oz. in 2 gals. of water. Apply as needed, but not more than 12 applications. Can be applied up to 7 days prior to harvest.
peppers	vegetable leafminers, cabbage looper, flea beetle, corn earworm, pepper weevil, European corn borer, green cloverworms, leafhoppers	Use ½ fl.oz. in 1 gal. of water. Apply as needed, but not more than 8 applications per season. Can be applied up to 3 days prior to harvest.
strawberries, raspberries, blueberries	strawberry weevils, aphids, spider mites, lygus bugs, flea beetles, omnivorous leafhopper, spittlebugs, Japanese beetles, field crickets, red spider, leafroller, weevils	Use ½ fl.oz. in 1 gal. water. Allow 14 days between last application and harvest. Make no more than 8 foliar applications per growing season.
sweet corn	armyworms (fall, southern), corn earworms, european corn borers	Use ¾ fl.oz. in 2 gals. of water. Apply every 5 days, but not more than 6 applications per season. Can be applied up to 1 day prior to harvest.
tomatoes	beet and southern armyworm, tomato pinworm, tomato fruitworm, horn worms, granulate cutworms, Colorado potato beetle, cabbage looper	Use ½ fl.oz. in 1 gal. water. Apply to ensure uniform coverage. Can be applied up to day of harvest.

NUISANCE PESTS IN OUTDOOR AREAS

GENERAL DIRECTIONS

Use 1 2/3 fl.ozs. in 1 quart of water, treating 400 linear feet with a 6-inch spray band.

OUTSIDE SURFACES OF BUILDING

Use Permethrin 10 E.W. on ants, clover mites, crickets, earwigs, millipedes, sowbugs (pillbugs) and spiders to help prevent infestation of buildings, treat the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Also, apply as a residual spray to outside surfaces of building, including porches, screens, window frames, eaves, patios, garages, refuse dumps and other areas where these pests congregate or have been seen.

OTHER OUTDOOR AREAS

Use Permethrin 10 E.W. on ants, crickets, earwigs, fleas, ticks (including ticks that may carry Lyme disease) for treatment of localized infestations of these insects in areas where there are weeds or bushy non-crop areas, spray infested areas thoroughly. For ants, thoroughly wet hills and runways. Repeat application only as needed to maintain insect control.

To reduce annoyance from: cluster flies, flying moths, gnats, house flies, mosquitoes. Spray outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages, around shrubbery, and other places where these insects alight or congregate.

DIRECTIONS FOR APPLYING (continued)

WOOD-INFESTING INSECTS (including termites*): for beetles (furniture, deathwatch, false powderpost, longhorned, powderpost), carpenter ants (and other wood-infesting ants), carpenter bees and termites (including dry-wood termites): Treat outside surfaces of buildings, porches, wooden decks and patios, wooden fences, window frames, doorways, foundations, eaves, patios, garages, garbage areas and other areas where you may find these pests.

Mix 2½ fl.oz. with 1 gallon of water to treat 100 sq.ft. Spray infested surfaces until thoroughly wet, avoiding dripping and runoff. Apply evenly on wood surfaces. To treat small areas, apply by brushing or spraying the diluted spray. For large areas, use a coarse, low pressure spray. Do not spray into electrical outlets or utilities.

Spray infested areas and areas likely to be infested, such as wood surfaces, voids and channels in damaged wood, spaces between wooden parts of a structure, and between wood and foundations where infestation is likely to occur, also around fenceposts, wood piles (do not use on firewood) and tree stumps.

Suitable protective clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of eye protection, protective head covering, chemical resistant gloves and footwear, a long-sleeved shirt and long pants or coveralls are recommended. Do not permit contact with treated surfaces until spray has dried. Treated confined areas, such as detached garages, under porches and decks, should be ventilated during and immediately following application before the areas are re-occupied.

For carpenter ants, spray around doors and windows and other places where ants may enter or crawl and hide. Also treat cracks and crevices or openings into wall voids or other areas where these ants may be present. For carpenter bees, spray or inject into holes and tunnels already made by the bees with a tank sprayer, and spray areas where they are beginning to build tunnels into the wood. For Termites* (including dry-wood termites): treat as for other wood-infesting insects. This product will kill workers and winged reproductive forms of termites in the treated areas.

*The purpose of such applications of ProCare Insecticide & Termiticide Concentrate 10% Permethrin for termites is to kill workers or winged reproductive forms which may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment, or foundation treatment but are merely a supplement. This product is not recommended as sole protection against termites. For active indoor termite infestations, get a professional inspection.

SOIL TERMITE TREATMENT DIRECTIONS

Subterranean Termites: DO NOT USE INSIDE DWELLINGS. For use around existing structures, not for pre-construction use. For post-construction treatment around residential buildings. Mix 6.7 oz of this product per 1 gallon of water (makes 0.5% water emulsion). Apply as specified below. One treatment should last up to 5 years if treated soil is left undisturbed. Kills and controls subterranean termites. Chemicals for soil treatment are used to establish a barrier against termites. The chemical emulsion must be adequately dispersed in the soil to provide a barrier between the wood in the structure and the termite colonies in the soil. Barriers may be established in the soil on the outside perimeter of the house by trenching using a rate of 4 gallons of emulsion per 10 linear feet per foot of depth. A complete treatment may require treatment with another EPA registered product to the inside perimeter of the foundation and to other crucial areas. For these situations, contact a licensed pest control operator for treatment. Do not use under slabs, in crawl spaces, or indoors. If termites are found within the house or structure, contact a licensed pest control operator for treatment.

Suitable protective clothing should be worn in confined or overhead areas to prevent exposure to eyes and skin. The use of eye protection, protective head covering, chemical resistant gloves and footwear, a long-sleeved shirt and long pants or coveralls are recommended. Do not permit contact with treated surfaces until spray has dried. Treated confined areas, such as detached garages, under porches and decks, should be ventilated during and immediately following application before the areas are re-occupied.

Remove all wood debris and wood forms. Around piers, pipes, chimney bases and along foundations of poured concrete, dig a trench 6 inches wide and 4 inches deep. For brick and block foundations, dig the trench about 12 inches deep. If the footing is more than 12 inches deep, make holes with a crowbar, pipe or rod about 1 foot apart that extend from the trench bottom to the top of the footing. Make holes closer in hard-packed clay soils. NEVER DIG BELOW THE TOP OF THE FOOTING. Apply the prepared emulsion (or solution) in the trench and to the backfill at the rate of 4 gallons per 10 linear feet per foot of depth. Cover the treated soil with a thin layer of untreated soil.

For raised porches, terraces and entrance slabs outdoors, drill holes at one foot intervals, 6 inches from the foundation, and pour one-half gallon of emulsion (or solution) into each hole. Refill holes.

Be sure to treat the interiors of accessible hollow block foundations at the same rate as above. Do not use in structural voids insulated with rigid foam. Be careful to have adequate ventilation when treating areas under structure. For severe infestations or treatment under slabs with heat ducts or radiating heat pipes, in basements and buildings with insufficient crawl space, etc., contact a licensed pest control operator or termite control service.

4.75"

4.75"

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages is forty (40).**

INSIDE TEXT (Even Pages)

4.25"

DIRECTIONS FOR APPLYING (continued)

NOTE: Avoid contamination of public and private water supplies by following these precautions. Do not allow a hose or any type of faucet extension to reach into the termite chemical solution while filling or mixing this material with water. Use of anti-back flow equipment will help ensure that the user of this chemical will not contaminate domestic water supplies. Refer to FHA (Federal Housing Administration), state and local specifications for safe distances of treatment areas from wells. Do not treat structures that have cisterns, wells, French drains, or sump systems. Soil in the vicinity of wells, cisterns or ponds should not be treated. For those situations, contact the local Cooperative Extension Service or consult a commercial pest control service. Do not treat soil that is water saturated or frozen or in any conditions where runoff or movement from the treatment area (site) is likely to occur.

When to reapply: Reapplication for subterranean termites should only be made when there has been a disruption of the chemical barrier in the soil due to a construction, excavation, landscaping, etc. Reapplication should be made as a spot treatment to those areas. Annual reapplication of the entire premises must be avoided.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle an open container in a manner as to prevent spillage.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Non-refillable container. Do not re-use or re-fill this container. Offer for recycling if available. Completely empty container. Dispose of container at an approved waste disposal facility or by incineration if allowed by state and local authorities. Do not use container in connection with food, feed, or drinking water.

Spills: If the container is leaking or material is spilled, absorb liquid with an absorbent such as kitty litter and carefully sweep up and wash exposed area with detergent and water. In spill or leak incidents, keep unauthorized people away.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS Keep Out of Reach of Children

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

FIRST AID

If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the Poison Control Center 800-222-1222. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may call the National Pesticide Information Center at 1-800-858-7378, Monday through Friday from 7:30 AM to 3:30 PM Pacific Time. After 3:30 PM Pacific Time and on weekends, call your poison control center.

NOTE TO PHYSICIAN AND VETERINARIANS: Do not administer or instill milk, cream or other substances containing vegetable or animal fats, which will enhance absorption of lipophilic substances.

4.75"

INSIDE TEXT (Odd Pages)

4.50"

PRECAUTIONARY STATEMENTS (continued)

HAZARDS TO HUMANS AND DOMESTIC ANIMALS Keep Out of Reach of Children

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic organisms including fish and invertebrates. Do not apply directly to water. Avoid contamination of ornamental fish ponds. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively, visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not use in undiluted form.

NOTE: Buyer assumes all risks of use, storage or handling of this product not in strict accordance with label DIRECTIONS and PRECAUTIONS.

WARRANTY STATEMENT

Gro Tec, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Gro Tec, Inc. To the extent consistent with applicable law, Gro Tec, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. In addition to the foregoing, no purchaser of this product (other than an end user) shall be entitled to any reimbursement for any loss suffered as a result of any suspension or cancellation of the registration for this product by the U.S. Environmental Protection Agency. Except, as expressly provided herein, Gro Tec, Inc. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, the exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Gro Tec, Inc. election, the replacement of this product.

NOTES

4.75"

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages is forty (40).**